

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked... The Appointment request is required, but make edits to the Interval and Duration as needed... Please add any additional orders on the last page*****

SJH AMB BLANK THERAPY PLAN

version 11/7/2023

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN CHEMO THERAPY INFUSION APPOINTMENT REQUEST			
☒	Infusion Appointment Request			

Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 30 minutes, Sched. Tolerance: Schedule appointment at most 0 days before or at most 0 days after

Appointment Request (continued)

Flush Line

		Interval	Defer Until	Duration
☒	SJH ONCBCN OP KVO			
☒	sodium chloride 0.9% (NS) infusion 250 mL, intravenous, at 30 mL/hr, Once, Starting when released KVO	Every visit	S	Until discontin'td
☒	dextrose 5% (D5W) infusion 250 mL, intravenous, at 30 mL/hr, Once, Starting when released KVO	Every visit	S	Until discontin'td

Hypersensitivity Medications

		Interval	Defer Until	Duration
☒	SJH ONCBCN HYPERSENSITIVITY MEDS THERAPY PLAN			
☒	IV/CVAD Line Care Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling.	PRN	S	Until discontin'td
☒	Deaccess/Discontinue CVAD No details available for preview	PRN	S	Until discontin'td
☒	Oxygen Therapy - Clinic Performed, Status: Future, Expected: S, Expires: S+396 Start for any grade of reaction. O2 Delivery Method: Nasal cannula Titrate to saturation of: 92% Indications of O2: Other (see comments)	PRN	S	Until discontin'td
☒	sodium chloride 0.9% (NS) infusion 1,000 mL, intravenous, at 500 mL/hr, Once as needed, Hypersensitivity reaction, Starting when released Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100. Start for any grade of reaction.	PRN	S	Until discontin'td
☒	diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.	PRN	S	Until discontin'td
☒	famotidine (PEPCID) injection 20 mg	PRN	S	Until discontin'td

	Interval	Defer Until	Duration
20 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction. ☒ dexamethasone (DECADRON) injection	PRN	S	Until discont'd
10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms ☒ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE injection 1 mg/mL	PRN	S	Until discont'd
0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released Do NOT administer via direct IV injection without dilution. Give for Grade III - SEVERE/Anaphylaxis Symptoms.			