

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked... The Appointment request is required, but make edits to the Interval and Duration as needed... Please add any additional orders on the last page*****

SJH DARBEPOETIN ALFA (ARANESP)

version 3/22/2023

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN INJECTION APPOINTMENT REQUEST			
☒	Injection Appointment Request	Every 2 weeks	S	6 treatments

Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 30 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after

Appointment Request (continued)

Labs

		Interval	Defer Until	Duration
☒	SJH ONCBCN LAB CBC AUTO DIFFERENTIAL (OR CBC 3 PART) LOCATION SPECIFIC RULE			
☐	CBC with Automated Diff	Every 2 weeks	S	Until discount'd
	No details available for preview Selection Condition: Lab location - SJH Oncology - Not London or Corbin			
☐	CBC Auto Diff 3 Parts	Every 2 weeks	S	Until discount'd
	No details available for preview Selection Condition: Lab location - SJH Oncology - Corbin			
☐	SJH ONCBCN LAB IRON STUDIES (CBC OR CBC 3 PART) LOCATION SPECIFIC RULE			
☒	Iron and TIBC			
	No details available for preview			
☒	Ferritin			
	No details available for preview			
☒	Reticulocyte count			
	No details available for preview			
☐	CBC with Automated Diff			
	No details available for preview Selection Condition: Lab location - SJH Oncology - Not London or Corbin			
☐	CBC Auto Diff 3 Parts			
	No details available for preview Selection Condition: Lab location - SJH Oncology - Corbin			
☒	Iron, serum			
	No details available for preview			
☒	Vitamin B12			
	No details available for preview			
☒	Folate			
	No details available for preview			

Medications

Medications (continued)

		Interval	Defer Until	Duration
☐	SJH ONCBCN THERAPY ARANESP 200 MCG			
☐	darbepoetin alfa-polysorbate (ARANESP) injection 200 mcg	Every 2 weeks	S	6 treatments
	200 mcg, subcutaneous, Once, Starting when released Review date of last iron studies and verify repeat orders are 12 weeks from last iron studies while on aranesp.			
☐	SJH ONCBCN THERAPY ARANESP 300 MCG			
☐	darbepoetin alfa-polysorbate (ARANESP) injection 300 mcg	Every 2 weeks	S	6 treatments
	300 mcg, subcutaneous, Once, Starting when released Review date of last iron studies and verify repeat orders are 12 weeks from last iron studies while on aranesp.			
☐	SJH ONCBCN THERAPY ARANESP 500 MCG			
☐	darbepoetin alfa-polysorbate (ARANESP) injection 500 mcg/ml	Every 2 weeks	S	6 treatments
	500 mcg, subcutaneous, Once, Starting when released Review date of last iron studies and verify repeat orders are 12 weeks from last iron studies while on aranesp.			

Hypersensitivity Medications

		Interval	Defer Until	Duration
☒	SJH ONCBCN HYPERSENSITIVITY MEDS THERAPY PLAN			
☒	IV/CVAD Line Care	PRN	S	Until discont'd
	Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling.			
☒	Deaccess/Discontinue CVAD	PRN	S	Until discont'd
	No details available for preview			
☒	Oxygen Therapy -	PRN	S	Until discont'd
	Clinic Performed, Status: Future, Expected: S, Expires: S+396 Start for any grade of reaction. O2 Delivery Method: Nasal cannula Titrate to saturation of: 92% Indications of O2: Other (see comments)			
☒	sodium chloride 0.9% (NS) infusion	PRN	S	Until discont'd

	Interval	Defer Until	Duration
1,000 mL, intravenous, at 500 mL/hr, Once as needed, Hypersensitivity reaction, Starting when released Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100.			
Start for any grade of reaction.			
☒ diphenhydrAMINE (BENADRYL) injection 50 mg	PRN	S	Until discont'd
50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒ famotidine (PEPCID) injection 20 mg	PRN	S	Until discont'd
20 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒ dexamethasone (DECADRON) injection	PRN	S	Until discont'd
10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms			
☒ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE injection 1 mg/mL	PRN	S	Until discont'd
0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released Do NOT administer via direct IV injection without dilution.			
Give for Grade III - SEVERE/Anaphylaxis Symptoms.			