

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked... The Appointment request is required, but make edits to the Interval and Duration as needed... Please add any additional orders on the last page*****

SJH DEGARELIX (FIRMAGON)

version 9/28/2022

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN INJECTION APPOINTMENT REQUEST			
☒	Injection Appointment Request	Every 28 days	S	13 treatments

Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 30 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after

Appointment Request (continued)

Labs

		Interval	Defer Until	Duration
☒	SJH ONCBCN LAB CBC AUTO DIFFERENTIAL (OR CBC 3 PART) LOCATION SPECIFIC RULE			
☐	CBC with Automated Diff No details available for preview Selection Condition: Lab location - SJH Oncology - Not London or Corbin	Every 28 days	S	13 treatments
☐	CBC Auto Diff 3 Parts No details available for preview Selection Condition: Lab location - SJH Oncology - Corbin	Every 28 days	S	13 treatments
☒	SJH ONCBCN LAB CMP (OR BMP/HF) W/ LOCATION SPECIFIC RULE			
☐	Comprehensive metabolic panel No details available for preview Selection Condition: Lab location - SJH Oncology - Not Blazer or Bob-O-Link	Every 28 days	S	13 treatments
☐	Oncology Basic Metabolic Panel No details available for preview Selection Condition: Lab location - SJH Oncology - Blazer	Every 28 days	S	13 treatments
☐	Hepatic function panel No details available for preview Selection Condition: Lab location - SJH Oncology - Bob-O-Link	Every 28 days	S	13 treatments
☒	SJH ONCBCN LABS PSA			
☐	PSA No details available for preview Selection Condition: Lab location - SJH Oncology - Mt. Sterling	Once	S	1 treatment
☐	Prostate Specific Antigen(SENDOUT) No details available for preview Selection Condition: Lab location - SJH Oncology - Flaget	Once	S	1 treatment

Medications

		Interval	Defer Until	Duration
☐	SJH ONCBCN THERAPY DEGARELIX (FIRMAGON) 240 MG			
☐	degarelix (FIRMAGON) injection 240 mg	Once	S	1 treatment

Medications (continued)

	Interval	Defer Until	Duration
240 mg, subcutaneous, Once, Starting at treatment start time			
☐ SJH ONCBCN THERAPY DEGARELIX (FIRMAGON) 80 MG			
☐ degarelix (FIRMAGON) injection 80 mg	Every 28 days	S+28	12 treatments
80 mg, subcutaneous, Once, Starting at treatment start time			

Hypersensitivity Medications

	Interval	Defer Until	Duration
☒ SJH ONCBCN HYPERSENSITIVITY MEDS THERAPY PLAN			
☒ IV/CVAD Line Care	PRN	S	Until discontin'td
Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling.			
☒ Deaccess/Discontinue CVAD	PRN	S	Until discontin'td
No details available for preview			
☒ Oxygen Therapy -	PRN	S	Until discontin'td
Clinic Performed, Status: Future, Expected: S, Expires: S+396 Start for any grade of reaction. O2 Delivery Method: Nasal cannula Titrate to saturation of: 92% Indications of O2: Other (see comments)			
☒ sodium chloride 0.9% (NS) infusion	PRN	S	Until discontin'td
1,000 mL, intravenous, at 500 mL/hr, Once as needed, Hypersensitivity reaction, Starting when released Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100.			
Start for any grade of reaction.			
☒ diphenhydrAMINE (BENADRYL) injection 50 mg	PRN	S	Until discontin'td
50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒ famotidine (PEPCID) injection 20 mg	PRN	S	Until discontin'td
20 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒ dexamethasone (DECADRON) injection	PRN	S	Until discontin'td
10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms			
☒ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE injection 1 mg/mL	PRN	S	Until discontin'td

	Interval	Defer Until	Duration
--	----------	-------------	----------

0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released
Do NOT administer via direct IV injection without dilution.

Give for Grade III - SEVERE/Anaphylaxis Symptoms.