

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked... The Appointment request is required, but make edits to the Interval and Duration as needed... Please add any additional orders on the last page*****

SJH MEROPENEM (MERREM)

version 5/7/2024

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN THERAPY INF APPOINTMENT REQUEST (60 MIN)			
☒	Infusion Appointment Request			

Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after

Appointment Request (continued)

Labs

	Interval	Defer Until	Duration
☒ SJH ONCBCN LAB CREATININE SERUM			
☒ Creatinine			
No details available for preview			
☒ SJH ONCBCN LAB CBC W/PLT COUNT AUTO DIFF (OR CBC 3 PART) LOCATION SPECIFIC RULE			
☐ CBC with Automated Diff			
No details available for preview			
Selection Condition: Lab location - SJH Oncology - Not London or Corbin			
☐ CBC Auto Diff 3 Parts			
No details available for preview			
Selection Condition: Lab location - SJH Oncology - Corbin			
☐ SJH ONCBCN LAB HEPATIC FUNCTION PANEL			
☐ Hepatic function panel	Every visit	S	Until discont'd
No details available for preview			

Nursing Communication

	Interval	Defer Until	Duration
☒ SJH ONCBCN NURSING COMMUNICATION CATHETER			
☒ Nursing Communication	Every visit	S	Until discont'd
Establish and/or maintain central venous catheter. Dressing changes weekly and PRN. Biopatch dressing PRN. May obtain blood from IV access for labs. May administer Cath-Flo (Activase) 2 mg for blood return.			

KVO Fluids

	Interval	Defer Until	Duration
☒ SJH ONCBCN OP KVO			
☒ sodium chloride 0.9% (NS) infusion	Every visit	S	Until discont'd
250 mL, intravenous, at 30 mL/hr, Once, Starting when released KVO			
☒ dextrose 5% (D5W) infusion	Every visit	S	Until discont'd

KVO Fluids (continued)

	Interval	Defer Until	Duration
250 mL, intravenous, at 30 mL/hr, Once, Starting when released KVO			

Medications

	Interval	Defer Until	Duration
☒ SJH ONCBCN THERAPY MEROPENEM 1 GM IVPB			
☒ meropenem (MERREM) 1 g in sodium chloride 0.9% (NS) 100 mL IVPB (CMPD)			
1 g, intravenous, Once, Starting when released			

Emergency Medications

	Interval	Defer Until	Duration
☐ SJH ONCBCN HYPERSENSITIVITY MEDS			
☐ IV/CVAD Line Care			
Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling.			
☐ Deaccess/Discontinue CVAD			
No details available for preview			
☐ Oxygen Therapy			
No details available for preview			
☐ sodium chloride 0.9% (NS) infusion			
1,000 mL, intravenous, at 500 mL/hr, Continuous PRN, Hypersensitivity reaction to chemotherapy, Starting when released, Until Discontinued Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100.			
Start for any grade of reaction.			
☐ diphenhydrAMINE (BENADRYL) injection 50 mg			
50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☐ famotidine (PEPCID) injection 20 mg			
20 mg, intravenous, Once as needed, Hypersensitivity reaction Start for any grade of reaction.			

Interval

Defer Until

Duration

- ☐ dexamethasone (DECADRON) injection

10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released
Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms

- ☐ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE
injection 1 mg/mL

0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released
Do NOT administer via direct IV injection without dilution.

Give for Grade III - SEVERE/Anaphylaxis Symptoms.