

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked...The Appointment request is required, but make edits to the Interval and Duration as needed...Please add any additional orders on the last page*****

SJH TEZPELUMAB-EKKO (TEZSPIRE)

version 3/6/2024

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN INJECTION APPOINTMENT REQUEST			
☒	Injection Appointment Request	Every 28 days	S	Until discont'd

Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 30 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after

Appointment Request (continued)

Medications

		Interval	Defer Until	Duration
☒	SJH ONCBCN MED TEZEPELUMAB-EKKO 210 MG SUB-Q			
☒	tezepelumab-ekko (TEZSPIRE) 210 mg/1.91 mL injection 210 mg	Every 28 days	S	Until discontin'td
	210 mg, subcutaneous, Once, Starting at treatment start time Administer as a SUBQ injection into the upper arm, thigh, or lower abdomen (avoiding areas within 2 inches of navel). Rotate injection sites; do not inject into areas where the skin is tender, bruised, erythematous, or hardened.			

Hypersensitivity Medications

		Interval	Defer Until	Duration
☒	SJH ONCBCN HYPERSENSITIVITY MEDS THERAPY PLAN			
☒	IV/CVAD Line Care	PRN	S	Until discontin'td
	Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling.			
☒	Deaccess/Discontinue CVAD	PRN	S	Until discontin'td
	No details available for preview			
☒	Oxygen Therapy -	PRN	S	Until discontin'td
	Clinic Performed, Status: Future, Expected: S, Expires: S+396 Start for any grade of reaction. O2 Delivery Method: Nasal cannula Titrate to saturation of: 92% Indications of O2: Other (see comments)			
☒	sodium chloride 0.9% (NS) infusion	PRN	S	Until discontin'td
	1,000 mL, intravenous, at 500 mL/hr, Once as needed, Hypersensitivity reaction, Starting when released Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100.			
	Start for any grade of reaction.			
☒	diphenhydrAMINE (BENADRYL) injection 50 mg	PRN	S	Until discontin'td
	50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒	famotidine (PEPCID) injection 20 mg	PRN	S	Until discontin'td
	20 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒	dexamethasone (DECADRON) injection	PRN	S	Until discontin'td

	Interval	Defer Until	Duration
10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms ☒ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE PRN S Until discont'd injection 1 mg/mL 0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released Do NOT administer via direct IV injection without dilution. Give for Grade III - SEVERE/Anaphylaxis Symptoms.			