

Infusion Therapy Order

Patient Name: _____ Date of Birth: _____

Address: _____ Height: _____ Weight: _____

Phone: _____ Allergies: _____

Primary Diagnosis: _____ ICD-10 Code: _____

Requested Infusion Start Date: _____ Frequency: _____

Infusion Location: _____ Duration: _____

Insurance Authorization: _____

Prescriber Information:

Prescriber Name: _____ NPI: _____

Address: _____

Phone: _____ Fax: _____

Prescriber Signature: _____

*****Please use a single line and mark through any orders you do not want if it is pre-checked...The Appointment request is required, but make edits to the Interval and Duration as needed...Please add any additional orders on the last page*****

SJH IRON INTERCHANGE SUCROSE (VENOFER 300 MG)

version 4/20/2023

Appointment Request

		Interval	Defer Until	Duration
☒	SJH ONCBCN THERAPY INF APPOINTMENT REQUEST (120 MIN)			
☒	Infusion Appointment Request	Every 2 days	S	Until discont'd

Appointment Request (continued)

	Interval	Defer Until	Duration
Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 120 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after			
☒ SJH ONCBCN LAB APPOINTMENT REQUEST			
☒ Oncology Lab Appointment Request	Once	S+83	1 treatment
Status: Future, Expected: S Approximate, Expires: S+365, Sched. Duration: 15 minutes, Sched. Tolerance: Schedule appointment at most 2 days before or at most 2 days after			

Labs

	Interval	Defer Until	Duration
☒ SJH ONCBCN LAB IRON STUDIES (CBC OR CBC 3 PART) LOCATION SPECIFIC RULE			
☒ Iron and TIBC	Once	S+83	1 treatment
No details available for preview			
☒ Ferritin	Once	S+83	1 treatment
No details available for preview			
☒ Reticulocyte count	Once	S+83	1 treatment
No details available for preview			
☐ CBC with Automated Diff	Once	S+83	1 treatment
No details available for preview			
Selection Condition: Lab location - SJH Oncology - Not London or Corbin			
☐ CBC Auto Diff 3 Parts	Once	S+83	1 treatment
No details available for preview			
Selection Condition: Lab location - SJH Oncology - Corbin			
☒ Iron, serum	Once	S+83	1 treatment
No details available for preview			
☒ Vitamin B12	Once	S+83	1 treatment
No details available for preview			
☒ Folate	Once	S+83	1 treatment
No details available for preview			
☒ SJH ONCBCN LAB CMP OR (BMP, HF) LOCATION SPECIFIC RULE			
☐ Comprehensive metabolic panel	Once	S+83	1 treatment

Labs (continued)

	Interval	Defer Until	Duration
No details available for preview Selection Condition: Lab location - SJH Oncology - Not Blazer or Bob-O-Link			
☐ Oncology Basic Metabolic Panel	Once	S+83	1 treatment
No details available for preview Selection Condition: Lab location - SJH Oncology - Bob-O-Link			
☐ Hepatic function panel	Once	S+83	1 treatment
No details available for preview Selection Condition: Lab location - SJH Oncology - Blazer			

Nursing Communication

	Interval	Defer Until	Duration
☐ SJH ONCBCN NURSING COMMUNICATION IRON			
☐ Nursing Communication	Every visit	S	Until discont'd
Wait 30 minutes after premeds before administering the iron product.			

Flush Line

	Interval	Defer Until	Duration
☒ SJH ONCBCN SODIUM CHLORIDE 0.9% 50 ML/HR - KVO			
☒ sodium chloride 0.9% (NS) bolus	Every visit	S	Until discont'd
500 mL, intravenous, As needed, For primary line., Starting at treatment start time, Until Discontinued Infuse at 50 ml/hr for the duration of treatment. May increase or decrease rate as needed.			

Pre-Medications

	Interval	Defer Until	Duration
☐ SJH ONCBCN PREMED (APAP 650 PO / DIPH 25 IVP / FAMO 20 IVP / METHYLPRED 40 IVP)			
☐ acetaminophen (TYLENOL) tablet 650 mg	Every visit	S	Until discont'd
650 mg, oral, Once, Starting at treatment start time Recommended maximum dose of acetaminophen is 4000 mg from all sources in 24 hours			
☐ diphenhydramine (BENADRYL) injection 25 mg	Every visit	S	Until discont'd

Pre-Medications (continued)

	Interval	Defer Until	Duration
☐ 25 mg, intravenous, Once, Starting at treatment start time famotidine (PF) injection 20 mg	Every visit	S	Until discont'd
☐ 20 mg, intravenous, Once, Starting at treatment start time methylPREDNISolone (SOLU-MEDROL) injection 40 mg	Every visit	S	Until discont'd
40 mg, intravenous, Once, Starting at treatment start time			

Medications

	Interval	Defer Until	Duration
☐ iron sucrose (Venofer) 3 treatments			
☐ iron sucrose (VENOFER) 300 mg in sodium chloride 0.9% (NS) 250 mL IVPB	Every 2 days	S	3 treatments
300 mg, intravenous, Once, Starting 30 minutes after treatment start time *If insurance denial ferumoxytol: interchange with iron sucrose (Venofer) QOD x 3 doses.			
K90.9 - intestinal malabsorption, unspecified. D50.0 iron deficient anemia secondary blood loss.			
☐ iron sucrose (Venofer) 5 treatments			
☐ iron sucrose (VENOFER) 300 mg in sodium chloride 0.9% (NS) 250 mL IVPB	Every 2 days	S	5 treatments
300 mg, intravenous, Once, Starting 30 minutes after treatment start time *If insurance denial ferumoxytol: interchange with iron sucrose (Venofer) QOD x 3 doses.			
K90.9 - intestinal malabsorption, unspecified. D50.0 iron deficient anemia secondary blood loss.			
☐ iron sucrose (Venofer) 7 treatments			
☐ iron sucrose (VENOFER) 300 mg in sodium chloride 0.9% (NS) 250 mL IVPB	Every 2 days	S	7 treatments
300 mg, intravenous, Once, Starting 30 minutes after treatment start time *If insurance denial ferumoxytol: interchange with iron sucrose (Venofer) QOD x 3 doses.			
K90.9 - intestinal malabsorption, unspecified. D50.0 iron deficient anemia secondary blood loss.			

Hypersensitivity Medications

	Interval	Defer Until	Duration
☒ SJH ONCBCN HYPERSENSITIVITY MEDS			
☒ IV/CVAD Line Care	PRN	S	Until discont'd

	Interval	Defer Until	Duration
☒ Access existing central venous access device or insert peripheral IV as needed. Restart peripheral IV catheter (as needed) for any signs of redness, pain, swelling. Deaccess/Discontinue CVAD	PRN	S	Until discont'd
☒ No details available for preview Oxygen Therapy	PRN	S	Until discont'd
☒ No details available for preview sodium chloride 0.9% (NS) infusion	PRN	S	Until discont'd
1,000 mL, intravenous, at 500 mL/hr, Continuous PRN, Hypersensitivity reaction to chemotherapy, Starting when released, Until Discontinued Initiate infusion at 500 ml/hr. Titrate up to 999ml/hr to maintain SBP greater than 100.			
☒ Start for any grade of reaction. diphenhydrAMINE (BENADRYL) injection 50 mg	PRN	S	Until discont'd
50 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Start for any grade of reaction.			
☒ famotidine (PEPCID) injection 20 mg	PRN	S	Until discont'd
20 mg, intravenous, Once as needed, Hypersensitivity reaction Start for any grade of reaction.			
☒ dexamethasone (DECADRON) injection	PRN	S	Until discont'd
10 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released Give for Grade II - MODERATE symptoms or Grade III - SEVERE/Anaphylaxis Symptoms			
☒ EPINEPHrine (ADRENALIN) HYPERSENSITIVITY USE injection 1 mg/mL	PRN	S	Until discont'd
0.3 mg, intraMUSCULAR, Once as needed, Hypersensitivity reaction, Starting when released Do NOT administer via direct IV injection without dilution.			
Give for Grade III - SEVERE/Anaphylaxis Symptoms.			